

Abstract Reproduction Form

ABSTRACTS MUST BE RECEIVED BY THURSDAY, SEPTEMBER 6, 1984

THE AMERICAN SOCIETY OF HEMATOLOGY

26th Annual Meeting

Person to whom all correspondence and notification regarding this abstract should be addressed:

Name A. E. Felice, M.D., Ph.D.

Address Medical Research Service, V.A. Medical Ctr.
Augusta, GA 30901

Phone Number (404) 724-5116, ext. 2457

Mail to: National Office

American Society of Hematology

Abstract Division

6900 Grove Rd.

Thorofare, New Jersey 08086

DO NOT FOLD!

Indicate membership:

☒ American Society of Hematology

☐ Council on Thrombosis

Please read abstract carefully before mailing. An additional charge of \$15 will be assessed for all editorial corrections (consult accompanying Rules).

Check Appropriate Category

- ☐ Red Cell Structure & Function
- ☐ Anemias
- ☒ Hemoglobinopathies and Thalassemias
- ☐ Hematopoietic Cell Proliferation and Differentiation
- ☐ Hematopoietic Growth Factors
- ☐ Granulocytes and Monocytes
- ☐ Lymphocytes
- ☐ Bone Marrow Failure
- ☐ Molecular Oncology
- ☐ Leukemias and Myeloproliferative Disorders
- ☐ Lymphomas and Plasma Cell Dyscrasias
- ☐ Immunohematology
- ☐ Transfusion
- ☐ Coagulation and Thrombosis
- ☐ Platelets
- ☐ Bone Marrow Transplantation

The submission fee is \$15. Make checks payable to the American Society of Hematology.

Abstracts, of work that has been published or presented at a national meeting prior to December 1, 1984 will not be accepted for the program or, if accepted, will not be presented. If similar work has been published or presented the original features of this study should be outlined in the covering letter.

The material in this abstract will have been presented at a national meeting or published as a full paper prior to December 1, 1984.

Yes ☐ No ☒

If accepted for a poster session,

☒ I would present

☐ I would not present

HB SYNTHESIS OF HUMAN LEUKEMIA DERIVED CELL LINES. A. E. Felice*, E. Marino*, J. Kraus* and V.C. McKie* (intr. by E.C. Abraham). V.A. Medical Center and Depts. of Cell and Molecular Biology, Pathology and Pediatrics, Medical College of Georgia, Augusta, GA.

K562 is a pluripotent human leukemia cell line that spontaneously produces variable numbers of hemoglobinized progeny. The production of Hb by K562 cells is usually enhanced by Hemin and some other compounds. Hemin stimulates the production of the embryonic Hb Gower I (or $\epsilon_2\zeta_2$), Hb Gower II (or $\alpha_2\epsilon_2$), Hb Gower III (or $\epsilon_2\gamma_2$) and Hb Portland I (or $\zeta_2\gamma_2$) and the fetal Hb F (or $\alpha_2\gamma_2$) and Hb Bart's (or γ_4). The γ globins are a mixture of G_γ , $A_\gamma I$ and $A_\gamma T$ globins. The Hb composition varied among 22 experiments without regard to passage number, Hemin dose, proportion of serum or the cell density. It is likely that Hemin augments the expression of a pre-established, though unstable, program of Hb synthesis. When the cells produced predominantly embryonic globins [$(\epsilon + \zeta) > 70\%$], the ϵ/ζ and the $\Sigma\alpha/\Sigma$ non- α ratios were balanced. When the cells produced predominantly γ globins [$(\epsilon + \zeta) < 30\%$], the major Hb in culture was Hb Bart's due to severe deficiencies of α globin. The $\Sigma\alpha/\Sigma$ non- α ratio was severely imbalanced. Despite the diversity of γ globin levels, the ratio of G_γ and A_γ globin produced was constant ($G_\gamma = 56.5\%$). A new cell line FM-4 was obtained from the peripheral blood of a $4\frac{1}{2}$ yr old Black boy with Ph-ive AML. The patient was in terminal stage with multiple organ relapse 2 years after diagnosis. The FM-4 cells reacted moderately to strongly positive for myeloperoxidase, sudan black B, ANAE and CE spontaneously and after exposure to a number of compounds. FM-4 cells did not produce Hb but FM-4 conditioned medium inhibited the Hb production of K562 cells.

Member's signature _____

(each member may sponsor two abstracts)